ELSEVIER

European Journal of Pharmacology 284 (1995) 337–339

ejp

Author index

Akehira, K., Nakane, Y., Hioki, K. and Taniyama, K.
Site of action of galanin in the cholinergic transmission of guinea pig small intestine (284) 149
Anticevich, S.Z., Hughes, J.M., Black, J.L. and Armour, C.L.
Induction of human airway hyperresponsiveness by tumour necrosis factor-α (284) 221
Anwyl, R., see Wu, J. (284) R1
Aoyagi, T., see Majima, M. (284) 1
Armour, C.L., see Anticevich, S.Z. (284) 221

Ballejo, G., see Martins, S.L.R. (284) 265
Baron, B.M., see Kehne, J.H. (284) 109
Bartolome, J.V., Chang, K.-J. and Bartolome, M.B.
The inhibition of ornithine decarboxylase activity in developing rat tissues by central nervous system β-endorphin is mediated by μ-opioid receptors, but not by δ- or ϵ-opioid receptors (284) 43
Bartolome, M.B., see Bartolome, J.V. (284) 43
Beattie, D.T., see Escott, K.J. (284) 69
Benazzouz, A., Boraud, T., Dubédat, P., Boireau, A., Stutzmann, J.-M. and Gross, C.
Riluzole prevents MPTP-induced parkinsonism in the rhesus monkey: a pilot study (284) 299
Black, J.L., see Anticevich, S.Z. (284) 221
Boireau, A., see Benazzouz, A. (284) 299
Boraud, T., see Benazzouz, A. (284) 299
Bos, E., see Du, X.Y. (284) 119
Brain, S.D., see Escott, K.J. (284) 69
Brixius, K., see Kojda, G. (284) 315

Cazes, M., Provost, D., Versigny, A. and Cloarec, A.
In vivo pharmacological characterization of UP 269-6, a novel non-peptide angiotensin II receptor antagonist (284) 157
Cervo, L. and Samanin, R.
Presynaptic 5-HT_{1A} receptors mediate the effect of ipsapirone on punished responding in rats (284) 249
Chan, S.L.F., see Tsoli, E. (284) 199
Chang, K.-J., see Bartolome, J.V. (284) 43
Chang, K.C. and Hahn, K.H.
Is α-adrenoceptor blockade responsible for atropine flush? (284) 331
Chen, K. and Zimmerman, B.G.
Angiotensin II-mediated renal vasoconstriction amenable to α_1-adrenoceptor blockade (284) 281
Church, J., see Sawyer, D.C. (284) 309
Cloarec, A., see Cazes, M. (284) 157
Connor, H.E., see Escott, K.J. (284) 69

De Klippel, N., see Smolders, I. (284) 83
De Oliveira, R.B., see Martins, S.L.R. (284) 265
Du, X.Y., Schoemaker, R.G., Bos, E. and Saxena, P.R.
Characterization of the positive and negative inotropic effects of acetylcholine in the human myocardium (284) 119
Du, X.Y., see Van Gelderen, E.M. (284) 51
Dubédat, P., see Benazzouz, A. (284) 299
Dusting, G.J., see Ralevic, V. (284) 231

Ebinger, G., see Smolders, I. (284) 83
Eblen, F., see Löschmann, P.-A. (284) 129
Emoto, M., see Inoue, Y. (284) 77
Escott, K.J., Beattie, D.T., Connor, H.E. and Brain, S.D.
The modulation of the increase in rat facial skin blood flow observed after trigeminal ganglion stimulation (284) 69

Feng, Y., see Tokuyama, S. (284) 101

Gidener, S., LePard, K.J. and Stephens, Jr., R.L.
8-OH-DPAT stimulates gastric acid secretion through a vagal-independent, adrenal-mediated mechanism (284) 19
Gommans, J., Hijzen, T.H., Maes, R.A., Mos, J. and Olivier, B.
Discriminative stimulus properties of flesinoxan: effects of enantiomers, (*S*)-UH301 and WAY-100635 (284) 135
Graff, C.B., see Jackson, D.L. (284) 321
Griffin, G., see Pertwee, R.G. (284) 241
Gross, C., see Benazzouz, A. (284) 299

Hahn, K.H., see Chang, K.C. (284) 331
Hargreaves, K.M., see Jackson, D.L. (284) 321
Harrison, B.L., see Kehne, J.H. (284) 109
Hashitani, H. and Suzuki, H.
Electrical and mechanical responses produced by nerve stimulation in detrusor smooth muscle of the guinea-pig (284) 177
Hatta, K., Wolterink, G. and Van Ree, J.M.
Prolyl-leucyl-glycinamide, thyrotropin-releasing hormone and β-endorphin-(10–16), like antidepressants, antagonize melatonin-induced behavioural changes in rats (284) 327
Hehl, S., see Moser, C. (284) 35
Helme, R.D., see Ralevic, V. (284) 231
Hijzen, T.H., see Gommans, J. (284) 135
Himes, C.S., see Tang, A.H. (284) 13
Hioki, K., see Akehira, K. (284) 149
Hirose, H., Ide, K., Sasaki, T., Takahashi, R., Kobayashi, M., Ikemoto, F., Yano, M. and Nishikibe, M.
Corrigendum to 'The role of endothelin and nitric oxide in modulation of normal and spastic cerebral vascular tone in the dog' [Eur. J. Pharmacol. 277 (1995) 77–87] (284) 335
Ho, I.K., see Tokuyama, S. (284) 101
Huffman, J.W., see Pertwee, R.G. (284) 241
Hughes, J.M., see Anticevich, S.Z. (284) 221

Idänpään-Heikkilä, J.J., Rauhala, P. and Männistö, P.T.
μ- and δ-opioid receptor antisense oligodeoxynucleotides antagonize morphine-induced growth hormone secretion in rats (284) 227
Ide, K., see Hirose, H. (284) 335
Ikeda, Y., see Majima, M. (284) 1
Ikemoto, F., see Hirose, H. (284) 335
Inoue, H., see Inoue, Y. (284) 77
Inoue, Y., Emoto, M., Inoue, H., Kaku, K. and Kaneko, T.
Characterization of the binding sites for [^{3}H]glibenclamide in rat liver membranes (284) 77
Isaka, N., see Miyahara, M. (284) 25

Ito, M., see Miyahara, M. (284) 25
Itoh, H., see Miyahara, M. (284) 25

Jackson, D.L., Graff, C.B., Richardson, J.D. and Hargreaves, K.M.
Glutamate participates in the peripheral modulation of thermal hyperalgesia in rats (284) 321

Kaku, K., see Inoue, Y. (284) 77
Kaneko, T., see Inoue, Y. (284) 77
Katori, M., see Majima, M. (284) 1
Kehne, J.H., Baron, B.M., Harrison, B.L., McCloskey, T.C., Palfreyman, M.G., Poirot, M., Salituro, F.G., Siegel, B.W., Slone, A.L., Van Giersbergen, P.L.M. and White, H.S.
MDL 100,458 and MDL 102,288: two potent and selective glycine receptor antagonists with different functional profiles (284) 109
Khalil, Z., see Ralevic, V. (284) 231
Klockgether, T., see Löschmann, P.-A. (284) 129
Kobayashi, M., see Hirose, H. (284) 335
Kojda, G., Brixius, K., Kottenberg, K., Nix, P., Schlüter, K.-D., Piper, H.M. and Noack, E.
The new NO donor SPM3672 increases cGMP and improves contraction in rat cardiomyocytes and isolated heart (284) 315
Konishi, T., see Miyahara, M. (284) 25
Kottenberg, K., see Kojda, G. (284) 315
Kuribayashi, Y., see Majima, M. (284) 1

Lainton, J.A.H., see Pertwee, R.G. (284) 241
La Riccia, M., see Pontieri, F.E. (284) 205
LePard, K.J., see Gidener, S. (284) 19
Löschmann, P.-A., Eblen, F., Wüllner, U., Wachtel, H. and Klockgether, T.
Lamotrigine has no antiparkinsonian activity in rat models of Parkinson's disease (284) 129

Ma, Z.-Q., Violani, E., Villa, F., Picotti, G.B. and Maggi, A.
Estrogenic control of monoamine oxidase A activity in human neuroblastoma cells expressing physiological concentrations of estrogen receptor (284) 171
Maes, R.A., see Gommans, J. (284) 135
Maggi, A., see Ma, Z.-Q. (284) 171
Mainero, C., see Pontieri, F.E. (284) 205
Majima, M., Ikeda, Y., Kuribayashi, Y., Mizogami, S., Katori, M. and Aoyagi, T.
Ebelactone B, an inhibitor of urinary carboxypeptidase Y-like kininase, prevents the development of deoxycorticosterone acetate-salt hypertension in rats (284) 1
Männistö, P.T., see Idänpään-Heikkilä, J.J. (284) 227
Martins, S.L.R., De Oliveira, R.B. and Ballejo, G.
Rat duodenum nitrergic-induced relaxations are cGMP-independent and apamin-sensitive (284) 265
McCloskey, T.C., see Kehne, J.H. (284) 109
McLarnon, J.G., see Sawyer, D.C. (284) 309
Meijer, F., Ruijter, J.M., Van Delft, J.L. and Van Haeringen, N.J.
Nitric oxide induces vascular permeability changes in the guinea pig conjunctiva (284) 61
Michotte, Y., see Smolders, I. (284) 83
Misawa, M., see Suzuki, T. (284) 195
Miyahara, M., Ito, M., Itoh, H., Shiraishi, T., Isaka, N., Konishi, T. and Nakano, T.
Isoenzymes of cyclic nucleotide phosphodiesterase in the human aorta: characterization and the effects of E4021 (284) 25
Mizogami, S., see Majima, M. (284) 1
Morgan, N.G., see Tsoli, E. (284) 199
Mori, T., see Suzuki, T. (284) 195
Morishita, W., see Shew, T.R. (284) 93
Mos, J., see Gommans, J. (284) 135
Moser, C., Hehl, S. and Neumcke, B.
Inhibition of ATP-sensitive K^+ channels of mouse skeletal muscle by disopyramide (284) 35
Mue, S., see Tanabe, J.-i. (284) 257

Nagase, H., see Suzuki, T. (284) 195
Nakane, Y., see Akehira, K. (284) 149
Nakano, T., see Miyahara, M. (284) 25
Naraba, H., see Ueno, A. (284) 211
Narita, M. and Tseng, L.F.
Stimulation of spinal δ-opioid receptors in mice selectively enhances the attenuation of δ-opioid receptor-mediated antinociception by antisense oligodeoxynucleotide (284) 185
Neumcke, B., see Moser, C. (284) 35
Nishikibe, M., see Hirose, H. (284) 335
Nix, P., see Kojda, G. (284) 315
Noack, E., see Kojda, G. (284) 315

Oh-ishi, S., see Ueno, A. (284) 211
Ohuchi, K., see Tanabe, J.-i. (284) 257
Olivier, B., see Gommans, J. (284) 135
Orzi, F., see Pontieri, F.E. (284) 205

Palfreyman, M.G., see Kehne, J.H. (284) 109
Passarelli, F., see Pontieri, F.E. (284) 205
Penn, N.W.
Potentiation of morphine analgesic action in mice by β-carotene (284) 191
Pertwee, R.G., Griffin, G., Lainton, J.A.H. and Huffman, J.W.
Pharmacological characterization of three novel cannabinoid receptor agonists in the mouse isolated vas deferens (284) 241
Picotti, G.B., see Ma, Z.-Q. (284) 171
Pidoplichko, V.I., see Wilsch, V.W. (284) 141
Piper, H.M., see Kojda, G. (284) 315
Poirot, M., see Kehne, J.H. (284) 109
Pontieri, F.E., Mainero, C., La Riccia, M., Passarelli, F. and Orzi, F.
Functional correlates of repeated administration of cocaine and apomorphine in the rat (284) 205
Provost, D., see Cazes, M. (284) 157

Ralevic, V., Khalil, Z., Helme, R.D. and Dusting, G.J.
Role of nitric oxide in the actions of substance P and other mediators of inflammation in rat skin microvasculature (284) 231
Rashed, S.M.K. and Songu-Mize, E.
Regulation of Na^+-pump activity by dopamine in rat tail arteries (284) 289
Rauhala, P., see Idänpään-Heikkilä, J.J. (284) 227
Reymann, K.G., see Wilsch, V.W. (284) 141
Richardson, J.D., see Jackson, D.L. (284) 321
Rowan, M.J., see Wu, J. (284) R1
Ruijter, J.M., see Meijer, F. (284) 61

Salituro, F.G., see Kehne, J.H. (284) 109
Samanin, R., see Cervo, L. (284) 249
Sarre, S., see Smolders, I. (284) 83
Sasaki, T., see Hirose, H. (284) 335
Sastry, B.R., see Shew, T.R. (284) 93
Sawyer, D.C., McLarnon, J.G. and Church, J.
(−)-β-Cyclazocine is an antagonist of NMDA receptor-mediated responses and a potent neuroprotectant in rat cortical neurons (284) 309
Saxena, P.R., see Du, X.Y. (284) 119
Saxena, P.R., see Van Gelderen, E.M. (284) 51
Schlüter, K.-D., see Kojda, G. (284) 315
Schoemaker, R.G., see Du, X.Y. (284) 119
Schoemaker, R.G., see Van Gelderen, E.M. (284) 51

Shew, T.R., Morishita, W. and Sastry, B.R.
Presynaptic actions of glutamate receptor agonists in the CA1 region of rat hippocampus in vitro (284) 93
Shiraishi, T., see Miyahara, M. (284) 25
Siegel, B.W., see Kehne, J.H. (284) 109
Sim, M.K. and Soh, K.S.
Effects of des-Asp-angiotensin I on the electrically stimulated contraction of the rabbit pulmonary artery (284) 215
Slone, A.L., see Kehne, J.H. (284) 109
Smolders, I., De Klippel, N., Sarre, S., Ebinger, G. and Michotte, Y.
Tonic GABA-ergic modulation of striatal dopamine release studied by in vivo microdialysis in the freely moving rat (284) 83
Soh, K.S., see Sim, M.K. (284) 215
Songu-Mize, E., see Rashed, S.M.K. (284) 289
Stephens, Jr., R.L., see Gidener, S. (284) 19
Stutzmann, J.-M., see Benazzouz, A. (284) 299
Suzuki, H., see Hashitani, H. (284) 177
Suzuki, T., Mori, T., Tsuji, M., Misawa, M. and Nagase, H.
Discriminative stimulus properties of morphine mediated by μ_1-opioid receptors (284) 195

Takahashi, R., see Hirose, H. (284) 335
Tanabe, J.-i., Watanabe, M., Mue, S. and Ohuchi, K.
Dexamethasone inhibits the production of macrophage inflammatory protein 2 in the leukocytes in rat allergic inflammation (284) 257
Tang, A.H. and Himes, C.S.
Apomorphine produced more yawning in Sprague-Dawley rats than in F344 rats: a pharmacological study (284) 13
Taniyama, K., see Akehira, K. (284) 149
Tokumasu, T., see Ueno, A. (284) 211
Tokuyama, S., Feng, Y., Wakabayashi, H. and Ho, I.K.
Possible involvement of protein kinases in physical dependence on opioids: studies using protein kinase inhibitors, H-7 and H-8 (284) 101
Tseng, L.F., see Narita, M. (284) 185
Tsoli, E., Chan, S.L.F. and Morgan, N.G.
The imidazoline I_1 receptor agonist, moxonidine, inhibits insulin secretion from isolated rat islets of Langerhans (284) 199
Tsuji, M., see Suzuki, T. (284) 195

Ueno, A., Tokumasu, T., Naraba, H. and Oh-ishi, S.
Involvement of bradykinin in endotoxin-induced vascular permeability increase in the skin of rats (284) 211
Uhlén, S., see Wikberg-Matsson, A. (284) 271

Van Delft, J.L., see Meijer, F. (284) 61
Van Gelderen, E.M., Du, X.Y., Schoemaker, R.G. and Saxena, P.R.
Carotid blood flow distribution, haemodynamics and inotropic responses following calcitonin gene-related peptide in the pig (284) 51
Van Giersbergen, P.L.M., see Kehne, J.H. (284) 109
Van Haeringen, N.J., see Meijer, F. (284) 61
Van Ree, J.M., see Hatta, K. (284) 327
Versigny, A., see Cazes, M. (284) 157
Villa, F., see Ma, Z.-Q. (284) 171
Violani, E., see Ma, Z.-Q. (284) 171

Wachtel, H., see Löschmann, P.-A. (284) 129
Wakabayashi, H., see Tokuyama, S. (284) 101
Watanabe, M., see Tanabe, J.-i. (284) 257
White, H.S., see Kehne, J.H. (284) 109
Wikberg, J.E.S., see Wikberg-Matsson, A. (284) 271
Wikberg-Matsson, A., Wikberg, J.E.S. and Uhlén, S.
Identification of drugs subtype-selective for α_{2A}-, α_{2B}-, and α_{2C}-adrenoceptors in the pig cerebellum and kidney cortex (284) 271
Wilsch, V.W., Pidoplichko, V.I. and Reymann, K.G.
Preferential block of desensitizing AMPA receptor in hippocampal neurons by γ-D-glutamylaminomethylsulfonic acid (284) 141
Wolterink, G., see Hatta, K. (284) 327
Wu, J., Anwyl, R. and Rowan, M.J.
β-amyloid-(1–40) increases long-term potentiation in rat hippocampus in vitro (284) R1
Wüllner, U., see Löschmann, P.-A. (284) 129

Yano, M., see Hirose, H. (284) 335

Zimmerman, B.G., see Chen, K. (284) 281